


Leptophilic Dark Matter with Z' Interactions

Rebecca Leane

In collaboration with Nicole Bell, Yi Cai and Anibal
Medina

arXiv: 1407.3001



Motivation


- WIMP model of Dark Matter (DM) well motivated
- WIMPs becoming more constrained, but constraints are based on DM-hadron interactions

....perhaps DM does not interact this way?

- Indirect detection experiments observed excess in cosmic ray positron fraction, suggesting DM annihilates to leptonic final states


Model Setup

$$\begin{aligned} \mathcal{L} = & \mathcal{L}_{SM} - \frac{1}{4} Z'_{\mu\nu} Z'^{\mu\nu} - \frac{\epsilon}{2} Z'_{\mu\nu} B^{\mu\nu} + i\bar{\chi}\gamma_{\mu}\partial^{\mu}\chi \\ & + \bar{\chi}\gamma^{\mu}(g_{\chi}^V + g_{\chi}^A\gamma^5)\chi Z'_{\mu} + \bar{\ell}\gamma^{\mu}(g_{\ell}^V + g_{\ell}^A\gamma^5)\ell Z'_{\mu} \\ & - m_{\chi}\bar{\chi}\chi + \frac{1}{2}m_{Z'}^2 Z'_{\mu} Z'^{\mu}, \end{aligned}$$



$\Gamma_{\chi} \otimes \Gamma_{\ell}$	$\sigma(\chi\chi \rightarrow \bar{\ell}\ell)$	$\sigma(\chi N \rightarrow \chi N)$	Gauge invariant?
$V \otimes V$	s -wave	1 (1-loop)	Yes
$A \otimes V$	p -wave	v^2 (1-loop)	Yes
$V \otimes A$	s -wave	-	No
$A \otimes A$	p -wave	-	No


Direct Detection





$$\sigma_{VV} = \frac{\mu_N^2}{9\pi} \left[\frac{\alpha_{EM} Z}{\pi \Lambda^2} \log \left(\frac{m_\ell^2}{\mu^2} \right) \right]^2, \quad \mathcal{L}_{eff} = \frac{1}{\Lambda^2} (\bar{\chi} \Gamma_\chi \chi) (\bar{\ell} \Gamma_\ell \ell)$$

$$\sigma_{AV} = \frac{\mu_N^2 v_\chi^2}{9\pi} \left(1 + \frac{\mu_N^2}{2m_N^2} \right) \left[\frac{\alpha_{EM} Z}{\pi \Lambda^2} \log \left(\frac{m_\ell^2}{\mu^2} \right) \right]^2 \quad \Lambda = \frac{m_{Z'}}{\sqrt{g_\chi g_\ell}}$$

Vector-vector Z' couplings



LHC phenomenology







FIG. 9. Invariant mass for four muons (left) and transverse momentum p_T for leading in p_T muon (right) for $pp \rightarrow 4\mu$ in the SM and Z' model (with $m_{Z'} = 60$ GeV, $m_\chi = 10$ GeV, $g_\mu = g_\chi = 0.1$), at $\sqrt{s} = 14$ TeV and $\mathcal{L} = 300$ fb^{-1} . The peak in the four muon invariant mass spectrum is a reconstruction of the Z mass.





FIG. 10. Invariant mass for first and second leading muons in p_T (left) and third and fourth leading muons in p_T (right) for $pp \rightarrow 4\mu$ in the SM and Z' model (with $m_{Z'} = 60$ GeV, $m_\chi = 10$ GeV, $g_\mu = g_\chi = 0.1$), at $\sqrt{s} = 14$ TeV and $\mathcal{L} = 300$ fb^{-1} . The mass of the Z' can be seen clearly as the resonance at $m_{Z'} = 60$ GeV in the invariant mass spectrum M_{12} .





FIG. 11. Invariant mass for first and second leading muons in p_T (left) and transverse momentum p_T for p_T leading muon (right) both before cuts, for $pp \rightarrow 4\mu$ in the SM and Z' model (with $m_{Z'} = 150$ GeV, $m_\chi = 10$ GeV, $g_\mu = g_\chi = 0.19$), at $\sqrt{s} = 14$ TeV and $\mathcal{L} = 3000$ fb^{-1} .










FIG. 12. Invariant mass of third and fourth leading in p_T muons before cuts (left) and after cuts (right), for $pp \rightarrow 4\mu$ in the SM and Z' model (with $m_{Z'} = 150$ GeV, $m_\chi = 10$ GeV, $g_\mu = g_\chi = 0.19$), at $\sqrt{s} = 14$ TeV and $\mathcal{L} = 3000$ fb^{-1} .






Conclusions

- We have considered a leptophilic WIMP scenario in which DM does not couple to SM quarks at tree-level, and instead couples only to SM leptons, which is consistent with the many null DM results to date.
- Despite the loop-suppressed nature of this process for direct detection, the resulting bounds are strong. Vector-vector couplings almost completely ruled out, axial vector only resonant production of DM left.
- We placed constraints from a recent ATLAS search, as well as the future exclusion/discovery reach. Electron parameter space extremely constrained, muons only resonant production, taus much more open.
- Despite the absence of tree-level interactions with quarks, this leptophilic dark matter model is strongly constrained by results from nuclear recoil and hadron collider experiments.

BACK UP SLIDES

DM Relic Density



- Larger couplings = subdominant contribution to the relic density,
- Smaller couplings = overclose universe unless additional annihilation channels present
- The $Z'Z'$ channel is kinematically open only for Z' mass $<$ DM mass, while for Z' mass $>$ DM mass, the freeze-out is determined by annihilation to leptons.
- The annihilation cross section to leptons has an s-wave contribution when vector-like Z' coupling to DM, but proceeds via a velocity suppressed p-wave contribution with axial-vector bilinear.

(g-2) constraints

$$\Delta(g-2)_e \sim \frac{g_\ell^2}{6\pi^2} \frac{m_\ell^2}{m_{Z'}^2}$$

$$g_e \lesssim 0.3 \frac{m_{Z'}}{\text{GeV}},$$

$$g_\mu \lesssim 6 \times 10^{-3} \frac{m_{Z'}}{\text{GeV}},$$

$$g_\tau \lesssim \frac{m_{Z'}}{\text{GeV}}.$$

LEP-II constraints

- **Z' constraints:**

For Z' masses greater than 209 GeV, the largest center-of-mass energy at which LEP operated, the constraints are expressed in terms of four-fermion contact operators, known as the compositeness bounds

$$g_e \lesssim 0.044 \times m_{Z'} / (200 \text{ GeV})$$

- **Mono-photon constraints:**

For Z' mass > 30 GeV these constraints are stronger than LUX, but are comparable to the LEP Z' bounds. For masses outside of this range, LUX is more constraining.