DETECTING DARK MATTER IN PLANETS

REBECCA LEANE SLAC NATIONAL ACCELERATOR LABORATORY

U. STOCKHOLM, OKC МАҮ 18^{тн} 2021

BASED ON:

2010.00015 w/ JURI SMIRNOV 2104.02068 w/ TIM LINDEN

Outline

- New Search for Dark Matter in Exoplanets
 - Why exoplanets?
 - Calculating the signal
 - Detecting the signal
 - Dark Matter mass and cross section sensitivity
 - Outlook: what's needed next

• New Search and Analysis of Jupiter in Gamma Rays

Dark Matter in Exoplanets



Dark Matter in Exoplanets



Dark Matter in Exoplanets



Why Exoplanets?

Advantage 1: Exploding Research Program

First exoplanet discovery: 1992 Almost all exoplanets we now know: 2010+ Majority of known exoplanets: **last five years**



Many upcoming telescopes and searches!

James Webb Space Telescope (JWST) Transiting Exoplanets Survey Satellite (TESS) Rubin/LSST Roman/WFIRST Gaia Spacecraft Optical Gravitational Lensing Experiment (OGLE) Two Micron All Sky Survey (2MASS) Wide-field Infrared Survey Explorer (WISE) Thirty Meter Telescope (TMT) Extremely Large Telescope (ELT) Gaia Near Infra-Red (GaiaNIR) Large Ultraviolet Optical Infrared Surveyor (LUVOIR) Habitable Exoplanet Imaging Mission (HabEx) Origins Space Telescope (OST)

Ample motivation to consider **new ways** this exploding research area can be used to probe new physics.

Advantage 2: Statistics

Estimates predict around 300 billion exoplanets in our galaxy!

To date:

4,301 confirmed exoplanets 5,633 exoplanet candidates



x 10^11



x 10^11





x 10^11

One Jupiter :(

Billions of Exoplanets! :)

Advantage 3: Low temperatures

- Exoplanets can be very cold, as they do not undergo nuclear fusion
 - Low temperatures allow for a clearer signal over background for DM heating
- Low core temperatures in part prevent DM evaporation, providing new sensitivity to lighter (sub-GeV) DM



Advantage 4: SIZE

Neutron Star

+ Easier to detect than neutron stars

Jupiters and Brown Dwarfs

Exoplanet Search Targets



Earths + Super Earths: Mass: 0.001– 0.01 Mjup Radius: ~0.1 - 1 Rjup





Brown dwarfs: Mass: 13 – 75 Mjup Radius: ~1 Rjup Very dense!



Rogue Planets: Cold and all alone!

Most commonly Jupiter-sized up to brown dwarf sized

Calculating Dark Matter Exoplanet Heating

Calculating Exoplanet Temperatures

 Contributions to temperature from external heat (i.e. nearby stars), internal heat (e.g. from formation or burning processes), and dark matter:

$$\Gamma_{\rm heat}^{\rm tot} = \Gamma_{\rm heat}^{\rm ext} + \Gamma_{\rm heat}^{\rm int} + \Gamma_{\rm heat}^{\rm DM} = 4\pi R^2 \,\sigma_{\rm SB} \,T^4 \,\epsilon$$

- External heat: assume zero, means we need exoplanets either very far from their host, or not bound at all (rogue planets)
- Internal heat: determined by cooling rate over time, choose old exoplanets (e.g. 1-10 gigayears old) to minimize internal heat

Calculating Exoplanet Temperatures

 Contributions to temperature from external heat (i.e. nearby stars), internal heat (e.g. from formation or burning processes), and dark matter:

$$\Gamma_{\rm heat}^{\rm tot} = \Gamma_{\rm heat}^{\rm ext} + \Gamma_{\rm heat}^{\rm int} + \Gamma_{\rm heat}^{\rm DM} = 4\pi R^2 \,\sigma_{\rm SB} \,T^4 \,\epsilon$$

Heat power from DM:

• DM density throughout Galaxy:

$$\rho_{\chi}(r) = \frac{\rho_0}{(r/r_s)^{\gamma}(1 + (r/r_s))^{3-\gamma}}$$

- Relevant velocities:
 - DM halo velocity
 - Exoplanet escape velocity

 $v_{\rm esc}^2 = 2G_N M/R$

$$\Gamma_{\text{heat}}^{\text{DM}} = f \,\pi R^2 \rho_{\chi}(r) \, v_0 \left(1 + \frac{3}{2} \frac{v_{\text{esc}}^2}{v_d(r)^2}\right)$$

DM Heating vs Internal Heat



1 parsec = 3.26 light years

Telescope Sensitivity

- Use James Webb Space Telescope (planned launch Oct 2021)
- Infrared sensitivity (~0.5 28 microns)
- Has many instruments and filters, relevant choice for maximum sensitivity depends on peak wavelength





Signal with James Webb

- Can see many stars/planets at once
- Assume exoplanets radiate as a blackbody lacksquare
 - Assume peak of blackbody temperature sets the sensitivity limit
- Near-Infrared Imager and Slitless \bullet Spectrometer (NIRISS) for T > 500 K
- Mid-Infrared Instrument (MIRI) for \bullet T = 100 - 500 K

Won't need new dedicated searches; can piggyback

S/N = 10 in 10 ks

40

F480M

F430M

F444W

4.5

5.0

F380M

Search Challenges

Dust backgrounds: Rescatter some wavelengths, which can reduce intensity and shift spectrum peaks

Stars per pixel important, can outshine exoplanet signal

Optimal sensitivity is outside 0.1 kpc (about 1 degree off the plane)

Exoplanet temperatures vs sensitivity

35 Mjup – 55 Mjup

• NFW, gNFW, Burkert are DM profiles, shaded area is exoplanet mass range

 Sensitivity truncates at ~0.1kpc, due to stars per pixel, and dust scattering

Leane + Smirnov, PRL '21

Exoplanet masses vs sensitivity

Mjup – 14 Mjup

14 Mjup – 35 Mjup

Lower masses: DM heat > internal heat at all positions

Higher masses:

Strongest signal towards Galactic Center, local DM heating signal difficult to outperform internal heat

New DM Search with Exoplanets

- Mock distribution of exoplanets with masses 20 – 50 Jupiters, gNFW profile, with and without DM heating
- Exoplanets can be used to map the Galactic DM density, given sufficient telescope sensitivity
- Identify exoplanets via other methods (e.g. microlensing) first, follow up with James Webb

Prospects for these searches?

Planet	Radius (R_{jup})	Mass (M_{jup})	Distance	Orbit	Temp (No DM)	Temp (with DM)	Ref
Epsilon Eridani b	1.21	1.55	3 pc	3.4 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[84]
Epsilon Indi A b	1.17	3.25	3.7 pc	11.6 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[85]
Gliese 832 b	1.25	0.68	4.9 pc	3.6 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[86]
Gliese 849 b	1.23	1.0	8.8 pc	2.4 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[87]
Thestias	1.19	2.3	10 pc	1.6 au	$\lesssim 200 \text{ K}$	$\lesssim 650~{ m K}$	[88]
Lipperhey	1.16	3.9	12.5 pc	5.5 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[89]
HD 147513 b	1.22	1.21	12.8 pc	1.3 au	$\lesssim 200~{\rm K}$	$\lesssim 650~{ m K}$	[90]
Gamma Cephei b	1.2	1.85	13.5 pc	2.0 au	$\lesssim 200 \text{ K}$	$\lesssim 650~{ m K}$	[91]
Majriti	1.16	4.1	13.5 pc	2.5 au	$\sim 218~{\rm K}$	$\lesssim 650~{ m K}$	[92]
47 Ursae Majoris d	1.2	1.64	14 pc	11.6 au	$\lesssim 200~{\rm K}$	$\lesssim 650~{ m K}$	[93]
Taphao Thong	1.2	2.5	14 pc	2.1 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[93]
Gliese 777 b	1.21	1.54	15.9 pc	4.0 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[94]
Gliese 317 c	1.21	1.54	15.0 pc	25.0 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[95]
q ¹ Eridani b	1.23	0.94	17.5 pc	2.0 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[87]
HD 87883 b	1.21	1.54	18.4 pc	3.6 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[96]
ν^2 Canis Majoris c	1.24	0.87	19.9 pc	2.2 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[97]
Psi^1 Draconis B b	1.21	1.53	22.0 pc	4.4 au	$\lesssim 200 \ {\rm K}$	$\lesssim 650~{ m K}$	[98]
HD 70642 b	1.19	1.99	29.4 pc	3.3 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[99]
HD 29021 b	1.2	2.4	31 pc	2.3 au	$\lesssim 200~{\rm K}$	$\lesssim 650~{ m K}$	[100]
HD 117207 b	1.2	1.9	32.5 pc	4.1 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[101]
Xolotlan	1.2	0.9	34.0 pc	1.7 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[102]
НАТ-Р-11 с	1.2	1.6	38.0 pc	4.1 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[103]
HD 187123 c	1.2	2.0	46.0 pc	4.9 au	$\lesssim 200 \ {\rm K}$	$\lesssim 650~{ m K}$	[104]
HD 50499 b	1.2	1.6	46.3 pc	3.8 au	$\lesssim 200~{ m K}$	$\lesssim 650~{ m K}$	[101]
Dim	1.0	1.1	10.1	0.0	2000 IZ	C CEO V	[105]

- Many candidates already exist!
- Gaia may be able to see up to around 90,000 planets within 100 pc (local search)
- WFIRST/Roman expects to detect least several thousand exoplanets in the inner galaxy

Cross section sensitivity

Leane + Smirnov, PRL '21

Actions for successful discovery/exclusion

- Successful launch with JWST
- Large statistical sample obtained to overcome systematics
- Detailed simulations of atmosphere effects including DM
- Simulations of age/cooling curves of Jupiters + Dwarfs *including DM*

Complementary Search with new limits

Complementary Searches

Two regimes:

 1. DM annihilates to short-lived mediators
 → heats planets

2. DM annihilates to long-lived mediators → escapes planets!

Complementary Searches

Two regimes:

 1. DM annihilates to short-lived mediators
 → heats planets

2. DM annihilates to long-lived mediators → escapes planets!

Dark Matter in Jupiter

Dark Matter in Jupiter

Why Jupiter?

BIG Hot

Jupiter BIG Cold

Solar Comparison

Sun Long-Lived Mediator Limits

Leane, Ng, Beacom (PRD '17) Leane + HAWC Collaboration (PRD '18)

Jupiter

Cooler than the Sun: MeV-DM mass sensitivity! No one ever looked!

New Dark Matter Limits

Leane + Linden, '21

Summary: Jupiter

- First search for gamma rays from Jupiter
 - No robust evidence, tentative excess at low energies
 - Motivates follow up w/ MeV telescopes

- New DM search with Jovian gamma rays
 - Strong constraints w/ long-lived particles: light masses and small cross sections

Summary: DM heated exoplanets

- The exoplanet program is rapidly accelerating, lots of new surprises and discoveries inevitable
- Examined how exoplanets can be used to discover DM, due to overheating from captured DM
 - Old, cold Jupiters and brown dwarfs ideal
- Actionable discovery or exclusion searches with new infrared telescopes
 - Signal traces DM density in the Galaxy
- New sensitivity to DM parameter space: DM-proton scattering up to six orders of magnitude stronger than other limits
- Exciting opportunities soon to realize search, several telescopes may be informative, new infrared window to Inner Galaxy
 - Oct 2021 James Webb launch!

EXTRA SLIDES

Cross section sensitivity

First Jovian Gamma-Ray Measurement

Leane + Linden, '21

Galactic Center Population Signal

- Use all the neutron stars, all the brown dwarfs
- Large population in the Galactic Center, DM density high: large rate!
- Our new signal follows matter density: DM density * stellar density
 - DM Halo annihilation scales with DM density squared

Galactic Center Population Signal

- Detectability: compare with known gamma-ray data
 - Use Fermi and H.E.S.S. data for Galactic Center
 - No model assumptions on mediator, other than must escape
 - Brown dwarfs very large signal!

RKL, Linden, Mukhopadyay, Toro, 2021

New Limits w/ Brown Dwarfs and Neutron Stars

RKL, Linden, Mukhopadyay, Toro, 2021

DM Equilibrium and Evaporation

- For maximal rate, want DM scattering and annihilation to be in equilibrium
 - Find DM reaches equilibrium by 1-10 Gigayears
- Lower end of DM mass sensitivity will stop due to DM becoming too light and evaporating out of the planet
 - Using temperature profiles for different exoplanets, find minimum mass, condition to remain bound is:

$$E_{\rm DM}^{\rm kin} = \frac{3}{2}T(r) < \frac{G_N M(r)m_{\chi}}{2r}$$

 Evaporation occurs for ~4 MeV DM mass in brown dwarfs, ~30 MeV DM mass in Jupiters

Deviations: DM-overdensities

Deviations: Non-Blackbody Spectra

Atmosphere effects can cause deviations from a blackbody

$$B(\nu, T) = \frac{2\nu^3 \epsilon}{\exp\left(\frac{2\pi\nu}{k_b T}\right) - 1}$$

Local DM-Heated Exoplanet Search

- Local fluxes easier to detect, so lower normalization from emissivity isn't a severe penalty
- Allows use of more powerful filters: best JWST filter sensitivity is with higher temps (in this case, higher wavelength peaks)
- Local exoplanets with lower emissivities can extend local sensitivity to DM heating

DM scattering cross section sensitivity

$$f = \frac{C_{\rm cap}}{C_{\rm max}} = \sum_{N=1}^{\infty} f_N$$

$$f_N = p(N,\tau) \left[1 - \kappa \exp\left(-\frac{3\left(v_N^2 - v_{esc}^2\right)}{2v_d^2}\right) \right]$$

$$\kappa = \left(1 + \frac{3}{2}\frac{v_{\rm N}^2}{v_d^2}\right) \left(1 + \frac{3}{2}\frac{v_{\rm esc}^2}{v_d^2}\right)^{-1}$$

Here v_d is the velocity dispersion, $v_N = v_{\rm esc} (1 - \langle z \rangle \beta)^{-N/2}$ where the average scattering angle is $\langle z \rangle = 1/2$ [143], $\beta = 4m_{\chi}m_A/(m_{\chi} + m_A)^2$, and m_A is the mass of the target particle. The probability that the DM particle scatters N times is

$$p(N,\tau) = \frac{2}{\tau^2} \left(N_s + 1 - \frac{\Gamma(N_s + 2, \tau)}{N_s!} \right) \quad \tau = \frac{3}{2} \frac{\sigma}{\sigma_{\text{sat}}}$$

 $\sigma_{\rm sat} = \pi R^2 / N_{\rm SM}$

$$\sigma_{\chi A}^{\rm SD} = \sigma_{\chi N}^{\rm SD} \left(\frac{\mu(m_A)}{\mu(m_N)}\right)^2 \frac{4(J+1)}{3J} \left[a_p \langle S_p \rangle + a_n \langle S_n \rangle\right]^2 \tag{1}$$

$$\sigma_{\chi A}^{\rm SI} = \sigma_{\chi N}^{\rm SI} \left(\frac{\mu(m_A)}{\mu(m_N)}\right)^2 \left[Z + \frac{a_n}{a_p}(A - Z)\right]^2$$

AGE – COOLING CURVES

DARK MATTER IN CELESTIAL OBJECTS

Apollo mission data: rock content and heat flux

Earth

20,000 boreholes Future Martian drilled kilometers deep mission: more info into the ground, internal heat measured Jupiter

DM heat

anomaly?

Ganymede

Uranus

Impact on

Volcanoes?

magnetic fields?

DM limits from temperature

Neutron Stars

DM heating, infrared telescopes

White Dwarfs

Sun

Neutrinos, long-lived particle decays outside the Sun

Evolution of dark matter number density

$$\frac{d}{dt}N_{\chi} = \Gamma_{\rm cap} - C_{\rm ann}N_{\chi}^2$$

LONG-LIVED SIGNAL BOOST:

Long-lived mediators

Schuster, Toro, Yavin (PRD '10) Batell, Pospelov, Ritz, Shang (PRD '10) Meade, Nussinov, Papucci, Volansky (JHEP '10)

LONG-LIVED SIGNAL BOOST: NEUTRINOS

LONG-LIVED SIGNAL BOOST: GAMMA RAYS

RL, Ng, Beacom (PRD '17)

NEW LIMITS WITH HAWC

HAWC Collaboration + **RL** (PRD '18) HAWC Collaboration + **RL** (PRD '18)

LONG-LIVED SIGNAL BOOST: GAMMA RAYS

RL, Ng, Beacom (PRD '17)

NEW LIMITS WITH FERMI AND HAWC

HAWC Collaboration + RL (PRD '18)

SOLAR DARK MATTER LIMITS

1

SOLAR DARK MATTER LIMITS: UPDATED

HAWC Collaboration + RL (PRD '18)

SOLAR DARK MATTER LIMITS: ALL

HAWC Collaboration + RL (PRD in press '18)

